

## Chapter 1: Systems of Linear Equations and Matrices

### Multiple Choice Questions

1. Which of the following equations is linear?

(A)  $2x_1^2 + 3x_2^3 + 4x_3^4 = 5$

(B)  $\sqrt{3}x_1 - \sqrt{2}x_2 + x_3 = 5$

(C)  $\sqrt{5}x_1 + 5\sqrt{x_2} - x_3 = 1$

(D)  $2^2x_1 + \cos(x_2) + 4x_3 = 7$

2. Which system corresponds to the following augmented matrix?

$$\left[ \begin{array}{cccc} 1 & 11 & 6 & 3 \\ 9 & 4 & 0 & -2 \end{array} \right]$$

(A)  $x_1 + 11x_2 = -3$   
 $9x_1 + 4x_2 = -2$

(B)  $x_1 + 11x_2 + 6x_3 = 3$   
 $9x_1 + 4x_2 = -2$

(C)  $x_1 + 11x_2 + 6x_3 + 3x_4 = 0$   
 $9x_1 + 4x_2 - 2x_4 = 0$

(D)  $x_1 + 9x_2 = 0$   
 $11x_1 + 4x_2 = 0$   
 $6x_1 = 0$   
 $3x_1 - 2x_2 = 0$

3. Which of the following statements best describes the following augmented matrix?

$$A = \left[ \begin{array}{cccc} 1 & 2 & 6 & 5 \\ -1 & 1 & -2 & 3 \\ 1 & -4 & -2 & 1 \end{array} \right]$$

(A)  $A$  is consistent with a unique solution.

(B)  $A$  is consistent with infinitely many solutions.

(C)  $A$  is inconsistent.

(D) none of the above.

4. Which of the following matrices is in *reduced* row echelon form?

$$(A) \begin{bmatrix} 1 & 0 & -1 & 1 \\ 0 & 1 & 2 & 0 \\ 0 & 1 & 3 & 1 \end{bmatrix}$$

$$(B) \begin{bmatrix} 1 & 0 & 2 & 5 \\ 0 & 1 & -7 & 5 \\ 0 & 0 & 1 & 14 \end{bmatrix}$$

$$(C) \begin{bmatrix} 1 & 0 & 0 & 11 & -3 \\ 0 & 0 & 0 & 1 & 4 \end{bmatrix}$$

$$(D) \begin{bmatrix} 1 & 0 & -5 \\ 0 & 1 & 3 \\ 0 & 0 & 0 \end{bmatrix}$$

5. If the matrix  $A$  is  $4 \times 2$ ,  $B$  is  $3 \times 4$ ,  $C$  is  $2 \times 4$ ,  $D$  is  $4 \times 3$ , and  $E$  is  $2 \times 5$ , which of the following expressions is *not* defined?

$$(A) A^T D + C B^T \quad (B) (B + D^T)A \quad (C) CA + C B^T \quad (D) DBAE$$

6. What is the second row of the product  $AB$ ?

$$A = \begin{bmatrix} 0 & 2 & 3 \\ 5 & 4 & 8 \\ 9 & 7 & 2 \end{bmatrix}, \quad B = \begin{bmatrix} 2 & 1 & 7 \\ 6 & 3 & 2 \\ 2 & 9 & 7 \end{bmatrix}$$

$$(A) \begin{bmatrix} 18 & 33 & 25 \end{bmatrix} \quad (B) \begin{bmatrix} 64 & 48 & 91 \end{bmatrix} \quad (C) \begin{bmatrix} 50 & 89 & 99 \end{bmatrix} \quad (D) \begin{bmatrix} 48 & 89 & 33 \end{bmatrix}$$

7. Which of the following is the determinant of the  $2 \times 2$  matrix  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ ?

$$(A) ad - bc \quad (B) bc - ad \quad (C) \frac{1}{bc - ad} \quad (D) \frac{1}{ad - bc}$$

8. Which of the following matrices is *not* invertible?

$$(A) \begin{bmatrix} 3 & 6 \\ 2 & 4 \end{bmatrix} \quad (B) \begin{bmatrix} 7 & 7 \\ 2 & 3 \end{bmatrix} \quad (C) \begin{bmatrix} 9 & 0 \\ 4 & 4 \end{bmatrix} \quad (D) \begin{bmatrix} 9 & 3 \\ 6 & 5 \end{bmatrix}$$

9. Which of the following matrices is *not* an elementary matrix?

$$(A) \begin{bmatrix} 1 & 0 \\ 5 & 1 \end{bmatrix} \quad (B) \begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix} \quad (C) \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (D) \begin{bmatrix} 1 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

10. For which elementary matrix  $E$  will the equation  $EA = B$  hold?

$$A = \begin{bmatrix} 1 & 4 & 6 \\ 0 & 0 & 1 \\ 2 & 10 & 9 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 4 & 6 \\ 0 & 0 & 1 \\ 0 & 2 & -3 \end{bmatrix}$$

$$(A) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} \quad (B) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \quad (C) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix} \quad (D) \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$$

11. Which matrix will be used as the inverted coefficient matrix when solving the following system?

$$3x_1 + x_2 = 4$$

$$5x_1 + 2x_2 = 7$$

$$(A) \begin{bmatrix} 2 & -1 \\ -5 & 3 \end{bmatrix} \quad (B) \begin{bmatrix} -2 & 1 \\ 5 & -3 \end{bmatrix} \quad (C) \begin{bmatrix} 2 & 1 \\ 5 & 3 \end{bmatrix} \quad (D) \begin{bmatrix} -2 & -1 \\ -5 & -3 \end{bmatrix}$$

12. What value of  $b$  makes the following system consistent?

$$4x_1 + 2x_2 = b$$

$$2x_1 + x_2 = 0$$

$$(A) b = -1 \quad (B) b = 0 \quad (C) b = 1 \quad (D) b = 2$$

13. If  $A$  is a  $3 \times 3$  diagonal matrix, which of the following matrices is *not* a possible value of  $A^k$  for some integer  $k$ ?

$$(A) \begin{bmatrix} 0 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 9 \end{bmatrix} \quad (B) \begin{bmatrix} 1 & 0 & 1 \\ 0 & 16 & 0 \\ 4 & 0 & 25 \end{bmatrix} \quad (C) \begin{bmatrix} 1 & 0 & 0 \\ 0 & \frac{1}{4} & 0 \\ 0 & 0 & -1 \end{bmatrix} \quad (D) \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

14. The matrix  $\begin{bmatrix} 3 & 0 & 0 \\ 0 & -7 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  is:

- (A) upper triangular.
- (B) lower triangular.
- (C) both (A) and (B).
- (D) neither (A) nor (B).

15. If  $A$  is a  $4 \times 5$  matrix, find the domain and codomain of the transformation  $T_A(\mathbf{x}) = A\mathbf{x}$ .

- (A) Not enough information
- (B) Domain:  $R^4$ , Codomain:  $R^5$
- (C) Domain:  $R^5$ , Codomain:  $R^5$
- (D) Domain:  $R^5$ , Codomain:  $R^4$

16. Which of the following is a matrix transformation?

- (A)  $T(x, y, z) = (yx^2, yz^2)$
- (B)  $T(x, y, z, w) = (xy, yz, zw, wx)$
- (C)  $T(x, y, z) = (x + 1, x + 2, x + z, y + z)$
- (D)  $T(x, y) = (4x, 5x, -x, 0)$

17. Which matrix represents reflection about the  $xy$ -plane?

- (A)  $\begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  (B)  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$  (C)  $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$  (D)  $\begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

18. Use matrix multiplication to find the image of the vector  $(2, 1)$  when it is rotated counterclockwise about the origin through an angle  $\theta = 45^\circ$ .

- (A)  $\left(\frac{\sqrt{2}}{2}, \frac{3\sqrt{2}}{2}\right)$  (B)  $\left(\frac{3\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$  (C)  $\left(-\frac{\sqrt{2}}{2}, \frac{3\sqrt{2}}{2}\right)$  (D)  $\left(-\frac{3\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\right)$

19. Which of the following pairs of operators  $T_1, T_2 : R^2 \rightarrow R^2$  commute? (That is, for which pair is it true that  $T_1 \circ T_2 = T_2 \circ T_1$ ?)

- (A)  $T_1$  is the reflection about the  $x$ -axis.  
 $T_2$  is the reflection about line  $y = x$ .
- (B)  $T_1$  is the orthogonal projection onto the  $x$ -axis.  
 $T_2$  is the reflection about line  $y = x$ .
- (C)  $T_1$  is the counterclockwise rotation about the origin through an angle of  $\pi$ .  
 $T_2$  is the projection onto the  $y$ -axis.
- (D)  $T_1$  is the reflection about the  $x$ -axis.  
 $T_2$  is the counterclockwise rotation about the origin through an angle of  $\pi/2$ .

### Free Response Questions

1. Find the relationship between  $a$  and  $b$  such that the following system has infinitely many solutions.

$$\begin{aligned} -x + 2y &= a \\ -3x + 6y &= b \end{aligned}$$

2. Solve the following system and use parametric equations to describe the solution set.

$$\begin{aligned}x_1 + 2x_2 + 3x_3 &= 11 \\2x_1 - x_2 + x_3 &= 2 \\3x_1 + x_2 + 4x_3 &= 13\end{aligned}$$

3. Determine whether the following system has no solution, exactly one solution, or infinitely many solutions.

$$\begin{aligned}2x_1 + 2x_2 &= 2 \\x_1 + x_2 &= 4\end{aligned}$$

4. Find the value of  $k$  that makes the system  $\begin{bmatrix} 15 & -3 & 6 \\ -10 & k & 9 \end{bmatrix}$  inconsistent.

5. Solve the following system using Gaussian elimination.

$$\begin{aligned}x_1 - x_2 - 5x_3 &= -1 \\-2x_1 + 2x_2 + 11x_3 &= 1 \\3x_1 - x_2 + x_3 &= 3\end{aligned}$$

6. Solve the following system for  $x$ ,  $y$ , and  $z$ .

$$\begin{aligned}\frac{1}{x} - \frac{1}{y} - \frac{1}{z} &= 0 \\ \frac{2}{x} + \frac{1}{y} + \frac{1}{z} &= 3 \\ \frac{3}{x} - \frac{1}{z} &= 0\end{aligned}$$

7. The curve  $y = ax^3 + bx^2 + x + c$  passes through the points  $(0, 0)$ ,  $(1, 1)$ , and  $(-1, -2)$ . Find and solve a system of linear equations to determine the values of  $a$ ,  $b$ , and  $c$ .

8. Solve the following system for  $x$  and  $y$ .

$$\begin{aligned}x^2 + y^2 &= 6 \\x^2 - y^2 &= 2\end{aligned}$$

9. Given  $C = \begin{bmatrix} 1 & -1 \\ 2 & 0 \end{bmatrix}$ , find  $CC^T$ .

10. Express the following matrix equation as a system of linear equations.

$$\begin{bmatrix} -1 & 7 & 0 \\ 0 & 4 & 3 \\ 6 & 0 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

11. Find the  $3 \times 3$  matrix  $A = [a_{ij}]$  whose entries satisfy the condition  $a_{ij} = i^2 - j$ .

12. Let  $A$  and  $B$  be  $n \times n$  matrices. Prove that  $\text{tr}(c \cdot A - B) = c \cdot \text{tr}(A) - \text{tr}(B)$ .

13. What is the inverse of  $\begin{bmatrix} 4 & 0 \\ 9 & 2 \end{bmatrix}$ ?

14. Given the polynomial  $p(x) = x^2 - 3x + 1$  and the matrix  $A = \begin{bmatrix} 4 & 4 \\ 6 & 1 \end{bmatrix}$ , compute  $p(A)$ .

15. Let  $A, B, C$ , and  $D$  be  $n \times n$  invertible matrices. Solve for  $A$  given that the following equation holds.

$$C^2DA^{-1}CB^{-1} = BCB^{-1}$$

16. Prove that for any  $m \times n$  matrices  $A$  and  $B$ ,  $(A - B)^T = A^T - B^T$ .

17. Use the inversion algorithm to find the inverse of the following matrix.

$$\begin{bmatrix} 1 & 2 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 4 \end{bmatrix}$$

18. Which elementary row operation will transform the following matrix into the identity matrix?

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -9 & 0 & 1 \end{bmatrix}$$

19. Find the  $3 \times 3$  elementary matrix that adds  $c$  times row 3 to row 1.

20. Find the elementary matrix  $E$  that satisfies

$$E \begin{bmatrix} 1 & 4 & 6 \\ 0 & 0 & 1 \\ 2 & 10 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 4 & 6 \\ 0 & 0 & 1 \\ 0 & 2 & -3 \end{bmatrix}$$

21. Solve the following system by inverting the coefficient matrix.

$$\begin{aligned} 7x + 2y &= 1 \\ 3x + y &= 5 \end{aligned}$$

22. Solve the following matrix equation for  $X$ .

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 4 \\ 5 & 6 & 0 \end{bmatrix} X = \begin{bmatrix} 2 & 2 & 3 & 0 \\ 0 & 0 & 0 & 1 \\ 3 & 1 & 1 & 1 \end{bmatrix}$$

23. Given that  $A^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 1 \\ 0 & 1 & 0 \end{bmatrix}$  and  $\mathbf{b} = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$ , solve the system  $A^2\mathbf{x} = \mathbf{b}$ .

24. Find a nonzero solution to the following equation.

$$\begin{bmatrix} 1 & 3 \\ 4 & -3 \end{bmatrix} \mathbf{x} = 3\mathbf{x}$$

25. Find the values of  $a$ ,  $b$ , and  $c$  that make the following matrix symmetric.

$$\begin{bmatrix} 3 & a & 2-b \\ 4 & 0 & a+b \\ 2 & c & 7 \end{bmatrix}$$

26. Let  $A = \begin{bmatrix} 3 & 4 & 3 \\ 0 & 0 & 6 \\ 0 & 0 & 2 \end{bmatrix}$ ,  $B = \begin{bmatrix} 1 & -7 & 6 \\ -4 & 5 & 0 \\ 1 & 0 & 2 \end{bmatrix}$ , and  $AB = [c_{ij}]$ .

Find the diagonal entries  $c_{11}$ ,  $c_{22}$ , and  $c_{33}$ .

27. Let the entries of a matrix  $A = [a_{ij}]$  be defined as  $a_{ij} = 2i^2 - i + j + g(j)$ , where  $g$  is a function of  $j$ . If  $A$  is a symmetric matrix, what is  $g(j)$ ?

28. Prove that for any square matrix  $A$ , the matrix  $B = (A + A^T)$  is symmetric.

29. Find the domain and codomain of the transformation defined by

$$\begin{bmatrix} 5 & 7 & 6 & 0 \\ 1 & 0 & -2 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}$$

30. Find the standard matrix for the operator  $T : R^2 \rightarrow R^2$  defined by

$$\begin{aligned} 3x_1 + x_2 &= w_1 \\ 4x_2 &= w_2 \end{aligned}$$

**31.** Find the standard matrix for the transformation  $T$  defined by the formula

$$T(x_1, x_2, x_3) = (x_1, -x_3, x_2 - x_1, 3x_2 + x_3)$$


**32.** Find the standard matrix  $A$  for the linear transformation  $T : R^2 \rightarrow R^2$  for which

$$T\left(\begin{bmatrix} 2 \\ -1 \end{bmatrix}\right) = \left(\begin{bmatrix} 3 \\ -4 \end{bmatrix}\right), T\left(\begin{bmatrix} -1 \\ 1 \end{bmatrix}\right) = \left(\begin{bmatrix} 5 \\ 2 \end{bmatrix}\right)$$

**33.** Prove that the composition of two rotation operators about the origin of  $R^2$  is another rotation about the origin.

**34.** Prove that if  $T_A : R^3 \rightarrow R^3$  and  $T_A(\mathbf{x}) = \mathbf{0}$  for every vector  $\mathbf{x}$  in  $R^3$ , then  $A$  is the  $3 \times 3$  zero matrix.

**35.** Write a balanced equation for the following chemical reaction.



**36.** Find the quadratic polynomial whose graph passes through the points  $(0, 3)$ ,  $(-1, 8)$ , and  $(1, 0)$ .

**37.** Use matrix inversion to find the production vector  $\mathbf{x}$  that meets the demand  $\mathbf{d}$  for the consumption matrix  $C$ .

$$C = \begin{bmatrix} 0.1 & 0.3 & 0.2 \\ 0.5 & 0.1 & 0.2 \\ 0.2 & 0.4 & 0.3 \end{bmatrix}; \mathbf{d} = \begin{bmatrix} 18 \\ 40 \\ 26 \end{bmatrix}$$



**Answers***Multiple Choice Answers*

1. (B)

2. (B)

3. (C)

4. (D)

5. (C)

6. (C)

7. (A)

8. (A)

9. (B)

10. (C)

11. (A)

12. (B)

13. (B)

14. (C)

15. (D)

16. (D)

17. (C)

18. (A)

19. (C)

*Free Response Answers*1.  $3a = b$

2.  $x_1 = -t + 3$ ,  $x_2 = -t + 4$ ,  $x_3 = t$

3. no solution

4.  $k = 2$

5.  $x_1 = 5$ ,  $x_2 = 11$ ,  $x_3 = -1$

6.  $x = 1$ ,  $y = -\frac{1}{2}$ ,  $z = \frac{1}{3}$

$$c = 0$$

7. System:  $a + b + c = 0$

$$-a + b + c = -1$$

Solution:  $a = \frac{1}{2}$ ,  $b = -\frac{1}{2}$ ,  $c = 0$

8.  $x = \pm 2$ ,  $y = \pm\sqrt{2}$

9.  $CC^T = \begin{bmatrix} 2 & 2 \\ 2 & 4 \end{bmatrix}$

$$-x + 7y = 0$$

10.  $4y + 3z = 0$

$$6x - 2z = 0$$

11.  $A = \begin{bmatrix} 0 & -1 & -2 \\ 3 & 2 & 1 \\ 8 & 7 & 6 \end{bmatrix}$

13.  $\begin{bmatrix} \frac{1}{4} & 0 \\ -\frac{9}{8} & \frac{1}{2} \end{bmatrix}$

14.  $\begin{bmatrix} 29 & 8 \\ 12 & 23 \end{bmatrix}$

15.  $A = B^{-1}C^2D$

17.  $\begin{bmatrix} 1 & -1 & \frac{1}{4} \\ 0 & \frac{1}{2} & -\frac{1}{4} \\ 0 & 0 & \frac{1}{4} \end{bmatrix}$

18. Add 9 times row 2 to row 4

$$19. \begin{bmatrix} 1 & 0 & c \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$20. E = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}$$

21.  $x = -9, y = 32$

$$22. \begin{bmatrix} -33 & -43 & -67 & 23 \\ 28 & 36 & 56 & -19 \\ -7 & -9 & -14 & 5 \end{bmatrix}$$

$$23. \mathbf{x} = \begin{bmatrix} 3 \\ 9 \\ 4 \end{bmatrix}$$

24. Any  $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$  such that  $2x_1 = 3x_2$ . Possible solution:  $\mathbf{x} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$

25.  $a = 4, b = 0, c = 4$

26.  $c_{11} = -10, c_{22} = 0,$  and  $c_{33} = 4$


27.  $g(j) = 2j^2 - 2j$

29. Domain:  $R^4$ , Codomain:  $R^2$

$$30. \begin{bmatrix} 3 & 1 \\ 0 & 4 \end{bmatrix}$$

$$31. \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ -1 & 1 & 0 \\ 0 & 3 & 1 \end{bmatrix}$$

$$32. \begin{bmatrix} 8 & 13 \\ -2 & 0 \end{bmatrix}$$



$$36. 3 - 4x + x^2$$

$$37. \mathbf{x} \approx \begin{bmatrix} 91.85 \\ 125.50 \\ 135.10 \end{bmatrix}$$